
All new STI episodes seen at GUM clinics in the UK: 1998 - 2007



Data is represented by country, gender and sexual orientation.

Notes on using these tables:

Numbers of diagnoses are adjusted for missing clinic data
Data are presented for the UK and England only.

Following recent ONS guidance on data disclosure, rules on the publication of STI data with small cell sizes have changed. Cells with values between 1 and 4 will now be anonymised with an asterix. In addition, where the anonymised cell can be deduced from the total, the next smallest cell size will also be anonymised. The only exception to this rule is for UK level data. Data for England and smaller geographic areas will now adopt the new policy.

Data Source:

England: GUM clinic KC60 returns
Wales: GUM clinic KC60 returns
Northern Ireland: GUM clinic KC60 returns
Scotland: GUM clinic ISD(D)5 returns from 31/03/2004 and STISS returns from 01/04/2004

Data Supplied by:

Health Protection Agency, Centre for Infections
Communicable Disease Surveillance Centre Wales
Communicable Disease Surveillance Centre Northern Ireland
Regional Information Branch, Department of Health, Social Services and Public Safety (DHSS&PS Northern Ireland)
ISD Scotland, Information Services, NHS National Services Scotland

Definitions of selected data totals:

Total new STI diagnoses: refer to 'Code Groups' for included KC60 conditions
Total other STI diagnoses: refer to 'Code Groups' for included KC60 conditions
Total other GUM clinic diagnoses: refer to 'Code Groups' for included KC60 conditions
Total services provided: refer to 'Code Groups' for included KC60 conditions

Data prepared by:

HIV and Sexually Transmitted Infections Department
Health Protection Agency
Centre for Infections
61 Colindale Avenue
London
NW9 5EQ

Tel. 020-8200-6868
Email: hiv/sti@hpa.org.uk
Website: www.hpa.org.uk

Data tables

United Kingdom

England

Code Groups

Further Information

England

Data on selected diagnoses, 1971 to 2007, is available upon request

HPA Centre for Infections

Contact: Geraldine Leong

Tel: +44 (0)20 8200 6868

Email: geraldine.leong@hpa.org.uk

Web: www.hpa.org.uk

Scotland

Data on selected diagnoses, 1956 to 2007, is available upon request

ISD Scotland

Contact: Kenny McIntyre

Tel: +44(0)131 275 6557;

Email: kenny.mcintyre@isd.csa.scot.nhs.uk

Web: www.isdscotland.org.uk

Wales

Data on selected diagnoses is available upon request.

National Public Health Service for Wales

Contact: Daniel Thomas

Tel: +44(0)29 2040 2472

email: daniel.thomas@nphs.wales.nhs.uk

Web: <http://www.nphs.wales.nhs.uk>

Northern Ireland

Data on selected diagnoses is available upon request.

Communicable Disease Surveillance Centre Northern Ireland

Contact: Brian Smyth

Tel: +44(0)2890 263 765

email: dbrian.smyth@hpa.org.uk

Web: www.hpa.org.uk



United Kingdom:

Number of new STI episodes seen at genitourinary medicine clinics by sex, 1998 - 2007

Diagnoses and/or treatment of infection or disease			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
A1,A2	Primary and secondary infectious syphilis	Male (all)	90	166	264	642	1,118	1,461	2,000	2,359	2,389	2,395
		-of which MSM	23	53	130	365	653	826	1,074	1,437	1,428	1,463
		Female	49	57	78	111	139	191	283	362	295	285
		Total	139	223	342	753	1,257	1,652	2,283	2,721	2,684	2,680
A3	Early latent syphilis (first 2 years)	Male (all)	91	127	134	275	363	459	571	711	753	906
		-of which MSM	20	31	51	118	194	211	301	424	466	542
		Female	58	79	93	104	129	156	201	189	196	163
		Total	149	206	227	379	492	615	772	900	949	1,069
A4,A5,A6	Other acquired syphilis	Male (all)	654	697	807	882	909	1,051	1,157	1,154	1,159	1,172
		-of which MSM	79	97	96	128	140	166	171	208	257	265
		Female	381	421	476	652	777	830	897	871	764	853
		Total	1,035	1,118	1,283	1,534	1,686	1,881	2,054	2,025	1,923	2,025
A7	Congenital syphilis, aged under 2	Male (all)	0	1	6	2	2	3	5	6	3	1
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	2	1	4	0	5	0	3	8	3	3
Total	2	2	10	2	7	3	8	14	6	4		
A8	Congenital syphilis, aged 2 or over	Male (all)	6	7	9	4	7	1	6	9	4	3
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	11	14	4	6	9	5	6	14	5	5
Total	17	21	13	10	16	6	12	23	9	8		
A9	Epidemiological treatment of suspected syphilis	Male (all)	20	31	27	64	168	270	423	535	700	777
		-of which MSM	0	7	6	25	86	161	255	324	451	509
		Female	36	35	26	30	52	70	95	126	133	165
Total	56	66	53	94	220	340	518	661	833	942		
B1,B2	Uncomplicated gonorrhoea	Male (all)	8,930	11,326	15,234	16,626	17,868	17,409	15,677	13,876	13,538	12,933
		-of which MSM	1,799	1,966	3,149	3,741	3,618	4,077	3,994	4,386	4,568	3,868
		Female	4,282	5,144	6,566	7,079	7,723	7,564	6,649	5,372	5,360	5,777
		Total	13,212	16,470	21,800	23,705	25,591	24,973	22,326	19,248	18,898	18,710
B3	Gonococcal ophthalmia neonatorum	Male (all)	3	6	7	7	8	10	10	12	21	13
		-of which MSM	-	-	-	-	-	-	-	-	-	
		Female	5	7	6	10	13	16	15	81	9	14
Total	8	13	13	17	21	26	25	93	30	27		
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	1,497	1,641	2,282	2,772	3,179	3,514	3,713	3,850	4,056	4,563
		-of which MSM	462	454	648	868	931	1,165	1,192	1,367	1,481	1,482
		Female	1,586	1,907	2,319	2,705	3,115	3,304	3,269	3,314	3,231	3,574
Total	3,083	3,548	4,601	5,477	6,294	6,818	6,982	7,164	7,287	8,137		
B5	Complicated gonococcal infection including PID and epididymitis	Male (all)	80	81	85	101	113	110	138	106	94	101
		-of which MSM	12	13	9	14	26	13	19	31	21	17
		Female	192	247	278	287	283	327	277	234	251	268
Total	272	328	363	388	396	437	415	340	345	369		

UK: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	46	34	32	30	38	65	88	262	221	262
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	27	24	33	36	39	83	85	67	80	124
		Total	73	58	65	66	77	148	173	329	301	386
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	21,064	24,435	29,818	33,362	38,676	43,188	48,625	51,911	56,187	60,798
		-of which MSM	504	690	1,082	1,531	1,646	2,113	2,228	2,712	3,358	3,690
		Female	27,662	32,556	38,514	43,153	48,912	52,971	56,114	57,507	57,596	61,188
		Total	48,726	56,991	68,332	76,515	87,588	96,159	104,739	109,418	113,783	121,986
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	368	480	569	610	632	699	724	810	845	878
		-of which MSM	31	26	25	40	35	54	49	74	69	71
		Female	1,680	1,959	2,365	2,411	2,462	2,426	2,617	2,532	2,480	2,809
		Total	2,048	2,439	2,934	3,021	3,094	3,125	3,341	3,342	3,325	3,687
C4D	Chlamydial ophthalmia neonatorum	Male (all)	15	18	20	15	18	23	49	22	17	12
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	17	13	29	29	26	26	41	23	23	8
		Total	32	31	49	44	44	49	90	45	40	20
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	10,305	12,823	15,700	19,471	23,941	25,173	27,420	29,305	28,948	32,335
		-of which MSM	213	277	479	726	1,147	1,491	1,613	1,913	2,219	2,562
		Female	7,732	9,221	11,202	13,291	15,673	16,317	18,691	20,709	20,972	23,055
		Total	18,037	22,044	26,902	32,762	39,614	41,490	46,111	50,014	49,920	55,390
C4H**	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	56,180	57,963	60,933	64,414	67,295	72,585	75,903	76,254	75,510	73,879
		-of which MSM	3,487	3,603	4,089	4,839	4,841	5,288	5,441	5,547	6,090	5,754
		Female	176	197	160	129	134	7,041	7,880	7,981	7,484	7,313
		Total	56,356	58,160	61,093	64,543	67,429	79,626	83,783	84,235	82,994	81,192
C4I	Epidemiological treatment of NSGI	Male (all)	5,496	6,413	6,497	6,363	6,287	6,441	6,598	6,130	6,086	6,489
		-of which MSM	461	733	646	723	670	800	829	775	869	827
		Female	19,361	19,880	20,410	21,161	20,122	18,236	18,613	18,323	17,391	17,383
		Total	24,857	26,293	26,907	27,524	26,409	24,677	25,211	24,453	23,477	23,872
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	2,717	3,341	3,865	4,045	4,214	4,280	4,675	5,427	5,083	5,593
		-of which MSM	159	221	216	248	239	257	286	368	354	377
		Female	10,295	10,790	11,431	11,534	11,964	11,386	13,243	12,721	13,110	14,843
		Total	13,012	14,131	15,296	15,579	16,178	15,666	17,918	18,148	18,193	20,436
C6A	Trichomoniasis	Male (all)	319	670	747	775	612	318	347	387	366	437
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	6,119	5,640	6,196	6,479	6,804	6,388	6,093	5,241	5,382	5,592
		Total	6,438	6,310	6,943	7,254	7,416	6,706	6,440	5,628	5,748	6,029
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	1,530	1,512	1,455	1,531	1,488	1,534	1,534	1,603	1,825	1,825
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	54,973	59,202	64,807	69,403	76,132	77,435	75,413	77,525	80,299	92,195
		Total	56,503	60,714	66,262	70,934	77,620	78,969	76,947	79,128	82,124	94,020
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	13,852	13,632	13,605	13,745	13,482	13,629	13,577	13,042	12,712	14,553
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	8,892	7,817	7,314	7,062	5,895	5,657	6,062	5,105	4,493	4,782
		Total	22,744	21,449	20,919	20,807	19,377	19,286	19,639	18,147	17,205	19,335

UK: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	
C7A	Anogenital candidosis	Male (all)	8,790	8,334	8,055	8,359	7,885	6,954	6,777	6,606	7,491	8,599	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	67,508	68,546	70,202	74,608	75,275	71,371	67,929	64,831	64,055	70,712	
		Total	76,298	76,880	78,257	82,967	83,160	78,325	74,706	71,437	71,546	79,311	
C7B	Epidemiological treatment of C6 & C7	Male (all)	3,694	3,688	3,386	3,761	4,276	4,332	4,285	4,438	4,227	4,804	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	6,289	6,468	5,807	6,704	6,741	7,133	8,004	7,386	7,297	8,200	
		Total	9,983	10,156	9,193	10,465	11,017	11,465	12,289	11,824	11,524	13,004	
C8,C9	Scabies/pediculosis pubis	Male (all)	5,132	5,044	5,225	4,687	4,091	3,524	2,799	2,230	2,183	2,067	
		-of which MSM	861	830	951	931	834	724	600	556	585	530	
		Female	1,654	1,452	1,296	1,062	825	593	456	341	298	303	
		Total	6,786	6,496	6,521	5,749	4,916	4,117	3,255	2,571	2,481	2,370	
C10A	Anogenital herpes simplex – first attack	Male (all)	6,688	6,660	6,812	7,200	7,218	7,209	7,237	7,596	8,460	10,031	
		-of which MSM	329	372	439	447	525	523	525	626	615	702	
		Female	10,560	10,849	11,011	11,744	12,220	12,024	11,836	12,234	13,337	16,031	
		Total	17,248	17,509	17,823	18,944	19,438	19,233	19,073	19,830	21,797	26,062	
C10B	Anogenital herpes simplex – recurrence	Male (all)	5,987	6,428	6,590	6,770	6,965	6,940	6,717	6,917	7,276	7,800	
		-of which MSM	382	422	463	438	440	503	510	568	589	645	
		Female	7,277	7,741	7,891	8,691	8,822	8,702	8,429	8,718	9,076	10,419	
		Total	13,264	14,169	14,481	15,461	15,787	15,642	15,146	15,635	16,352	18,219	
C11A	Anogenital warts – first attack	Male (all)	36,522	37,482	37,633	38,522	39,887	40,601	42,727	42,979	44,238	47,239	
		-of which MSM	1,672	1,760	1,877	1,987	2,231	2,397	2,396	2,593	2,667	2,841	
		Female	33,769	34,266	33,684	34,936	35,082	35,998	37,332	38,222	39,386	42,599	
		Total	70,291	71,748	71,317	73,458	74,969	76,599	80,059	81,201	83,624	89,838	
C11B	Anogenital warts – recurrence	Male (all)	25,462	25,385	25,420	25,620	24,957	24,718	25,189	26,433	27,935	30,681	
		-of which MSM	1,330	1,325	1,337	1,484	1,513	1,693	1,801	1,741	1,912	2,189	
		Female	13,994	13,861	14,051	13,869	13,742	14,188	14,622	15,631	17,025	19,106	
		Total	39,456	39,246	39,471	39,489	38,699	38,906	39,811	42,064	44,960	49,787	
C11C	Anogenital warts – re-registered cases	Male (all)	10,113	10,435	9,992	9,864	9,272	9,535	9,261	9,526	10,114	10,345	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	8,386	8,535	8,036	7,869	7,107	7,202	7,126	7,447	7,623	7,813	
		Total	18,499	18,970	18,028	17,733	16,379	16,737	16,387	16,973	17,737	18,158	
C12	Molluscum contagiosum	Male (all)	4,711	5,186	5,594	6,049	6,704	7,544	8,728	9,698	10,605	11,704	
		-of which MSM	228	238	221	243	231	273	285	325	351	374	
		Female	2,616	2,897	2,955	3,277	3,512	3,815	4,277	4,684	4,973	5,436	
		Total	7,327	8,083	8,549	9,326	10,216	11,359	13,005	14,382	15,578	17,140	
C13	Antigen positive viral hepatitis B	Male (all)	589	565	669	748	962	23	31	17	25	40	
		-of which MSM	112	109	135	165	203	9	15	6	3	15	
		Female	211	264	238	364	496	6	7	3	7	13	
		Total	800	829	907	1,112	1,458	29	38	20	32	53	
C13A**	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	0	0	0	0	0	619	672	655	653	697	
		-of which MSM	0	0	0	0	0	114	93	91	62	61	
		Female	0	0	0	0	0	276	328	329	344	359	
		Total	0	0	0	0	0	895	1,000	984	997	1,056	

UK: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
C13B**	**number of which were acute viral hepatitis B	Male (all)	0	0	0	0	0	47	55	48	43	44
		-of which MSM	0	0	0	0	0	14	19	14	8	8
		Female	0	0	0	0	0	11	12	30	26	19
		Total	0	0	0	0	0	58	67	78	69	63
C13C**	Viral hepatitis B: subsequent presentation	Male (all)	0	0	0	0	0	348	462	419	438	495
		-of which MSM	0	0	0	0	0	121	205	207	230	168
		Female	0	0	0	0	0	151	169	165	193	204
		Total	0	0	0	0	0	499	631	584	631	699
C14**	Viral hepatitis C: first diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	1,069	957	980	955	1,096	1,053	1,116	1,029	961	886
		-of which MSM	132	83	106	105	145	110	126	123	128	87
		Female	456	454	445	396	503	489	472	447	397	333
		Total	1,525	1,411	1,425	1,351	1,599	1,542	1,588	1,476	1,358	1,219
E1A	New HIV diagnosis - asymptomatic	Male (all)	1,166	1,125	1,244	1,713	2,365	2,737	2,922	3,154	2,963	3,633
		-of which MSM	585	551	624	770	772	1,134	1,299	1,501	1,411	1,609
		Female	374	589	691	1,137	2,012	2,154	2,116	2,110	1,815	2,316
		Total	1,540	1,714	1,935	2,850	4,377	4,891	5,038	5,264	4,778	5,949
E1B**	Asymptomatic HIV infection – subsequent presentation	Male (all)	7,000	7,502	8,173	8,758	10,387	198	629	832	915	980
		-of which MSM	490	557	122	76	47	84	426	656	691	775
		Female	1,964	2,280	3,006	3,872	5,630	56	222	202	307	312
		Total	8,964	9,782	11,179	12,630	16,017	254	851	1,034	1,222	1,292
E1BE2B**	Subsequent HIV presentation (not AIDS)	Male (all)	0	0	0	0	0	26,102	31,372	38,437	39,960	40,518
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	0	0	0	0	0	12,318	16,561	20,823	21,818	23,323
		Total	0	0	0	0	0	38,420	47,933	59,260	61,778	63,841
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	612	548	513	684	772	695	701	767	662	511
		-of which MSM	320	294	244	330	265	278	263	276	253	177
		Female	202	198	210	335	457	459	402	453	373	322
		Total	814	746	723	1,019	1,229	1,154	1,103	1,220	1,035	833
E2B**	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	8,590	8,689	9,421	9,508	9,597	58	433	502	467	481
		-of which MSM	508	521	104	7	8	38	328	405	357	350
		Female	1,902	2,065	2,376	2,699	3,381	8	107	79	76	98
		Total	10,492	10,754	11,797	12,207	12,978	66	540	581	543	579
E3A**	AIDS – first presentation	Male (all)	448	414	383	448	468	3	4	5	7	7
		-of which MSM	196	167	189	175	177	0	3	1	4	6
		Female	135	112	141	217	237	0	2	0	2	0
		Total	583	526	524	665	705	3	6	5	9	7
E3A1**	AIDS: first presentation - new HIV diagnosis	Male (all)	0	0	0	0	0	182	222	231	190	227
		-of which MSM	0	0	0	0	0	73	62	77	64	75
		Female	0	0	0	0	0	143	120	150	90	96
		Total	0	0	0	0	0	325	342	381	280	323
E3A2**	AIDS: first presentation - HIV diagnosed previously	Male (all)	0	0	0	0	0	240	156	162	160	113
		-of which MSM	0	0	0	0	0	97	76	61	82	54
		Female	0	0	0	0	0	135	65	70	70	46
		Total	0	0	0	0	0	375	221	232	230	159

UK: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	
E3B	AIDS – subsequent presentation	Male (all)	5,809	6,060	6,625	7,046	7,958	9,761	10,582	11,986	11,917	11,568	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	1,201	1,381	1,683	2,125	2,666	4,060	4,418	5,042	5,220	5,440	
Total			7,010	7,441	8,308	9,171	10,624	13,821	15,000	17,028	17,137	17,008	
D2A	Urinary tract infection	Male (all)	2,040	2,138	2,010	2,101	2,625	2,626	2,588	2,134	2,104	2,365	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	8,955	9,349	9,642	9,743	9,480	9,042	10,119	9,691	10,065	12,528	
Total			10,995	11,487	11,652	11,844	12,105	11,668	12,707	11,825	12,169	14,893	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	55,630	56,498	58,716	58,976	62,338	69,190	68,597	70,648	72,310	83,502	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	51,692	51,699	56,342	57,158	58,182	56,791	57,674	58,590	60,004	67,581	
Total			107,322	108,197	115,058	116,134	120,520	125,981	126,271	129,238	132,314	151,083	
D3	Other episodes not requiring treatment	Male (all)	108,395	112,394	118,230	139,176	158,614	151,051	183,482	207,164	229,372	254,921	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	116,237	120,252	123,430	143,377	161,215	150,515	176,853	204,369	224,070	256,795	
Total			224,632	232,646	241,660	282,553	319,829	301,566	360,335	411,533	453,442	511,716	
P1A	HIV antibody test (no sexual health screen)	Male (all)	77,413	79,819	90,280	117,896	156,922	58,621	57,271	42,381	36,429	33,271	
		-of which MSM	11,887	12,041	13,340	17,432	20,186	10,010	9,695	8,271	7,830	7,150	
		Female	61,758	63,346	71,831	98,559	138,304	44,666	41,445	31,737	27,209	25,401	
Total			139,171	143,165	162,111	216,455	295,226	103,287	98,716	74,118	63,638	58,672	
P1B	HIV antibody test offered and refused	Male (all)	13,484	13,825	15,220	20,396	45,408	100,363	127,092	127,295	122,586	117,323	
		-of which MSM	1,167	1,492	1,234	1,765	3,335	6,188	6,973	7,194	7,230	6,387	
		Female	12,481	13,086	14,236	20,035	50,304	114,577	143,218	145,051	141,750	146,336	
Total			25,965	26,911	29,456	40,431	95,712	214,940	270,310	272,346	264,336	263,659	
P2	Hepatitis B vaccination (1st dose only)	Male (all)	10,233	10,324	11,277	12,427	13,996	14,967	16,659	16,506	17,095	18,045	
		-of which MSM	4,967	4,892	5,387	6,114	6,961	8,095	8,779	8,969	8,945	9,149	
		Female	2,671	2,574	2,718	3,243	4,008	4,222	5,252	5,637	5,992	6,620	
Total			12,904	12,898	13,995	15,670	18,004	19,189	21,911	22,143	23,087	24,665	
P3	Contraception (excluding condom provision)	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	32,182	30,705	33,389	34,075	31,923	33,102	34,664	35,861	39,800	49,102	
Total			32,182	30,705	33,389	34,075	31,923	33,102	34,664	35,861	39,800	49,102	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	10,036	8,060	6,509	4,926	3,849	5,358	3,450	1,771	1,497	1,545	
Total			10,036	8,060	6,509	4,926	3,849	5,358	3,450	1,771	1,497	1,545	

UK: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1,412	1,231	1,146	1,033	766	924	813	398	304	277
		Total	1,412	1,231	1,146	1,033	766	924	813	398	304	277
S1**	Sexual health screen (no HIV antibody test)	Male (all)	0	0	0	0	0	151,572	153,409	147,782	141,924	142,310
		-of which MSM	0	0	0	0	0	9,952	10,083	11,329	12,605	12,349
		Female	0	0	0	0	0	185,799	184,799	180,119	172,421	179,404
		Total	0	0	0	0	0	337,371	338,208	327,901	314,345	321,714
S2**	HIV antibody test and sexual health screen	Male (all)	0	0	0	0	0	163,154	214,136	281,824	330,390	370,108
		-of which MSM	0	0	0	0	0	16,313	19,549	25,980	31,382	32,211
		Female	0	0	0	0	0	159,738	207,426	272,868	318,056	371,028
		Total	0	0	0	0	0	322,892	421,562	554,692	648,446	741,136
Total new STI diagnoses		Male (all)	144,625	154,541	168,567	179,460	191,602	202,606	213,513	218,047	223,534	232,688
		-of which MSM	10,010	10,617	13,056	15,486	15,916	18,030	18,521	20,509	21,835	21,548
		Female	99,657	106,865	115,468	123,709	132,568	143,562	149,776	150,211	152,309	165,302
		Total	244,282	261,406	284,035	303,169	324,170	346,168	363,289	368,258	375,843	397,990
Total other STI diagnoses		Male (all)	81,496	86,677	92,093	97,869	104,486	114,500	124,295	136,534	140,498	148,764
		-of which MSM	4,141	4,591	4,141	4,768	5,353	6,409	7,505	8,443	9,378	10,206
		Female	64,048	67,553	71,753	77,330	81,506	85,629	93,383	101,670	103,944	111,780
		Total	145,544	154,230	163,846	175,199	185,992	200,129	217,678	238,204	244,442	260,544
Total other GUM clinic diagnoses		Male (all)	87,194	87,324	88,877	90,176	94,153	100,356	99,694	100,639	102,789	117,810
		-of which MSM	244	192	241	270	348	368	458	441	431	339
		Female	210,424	213,090	222,452	231,398	237,319	234,645	230,456	226,271	228,982	258,748
		Total	297,618	300,414	311,329	321,574	331,472	335,001	330,150	326,910	331,771	376,558
Total services provided		Male (all)	209,525	216,362	235,007	289,895	374,940	639,728	752,049	822,952	877,796	935,978
		-of which MSM	18,021	18,425	19,961	25,311	30,482	50,558	55,079	61,743	67,992	67,246
		Female	225,329	229,963	245,604	299,289	385,754	692,619	793,657	875,642	929,298	1,034,686
		Total	434,854	446,325	480,611	589,184	760,694	1,332,347	1,545,706	1,698,594	1,807,094	1,970,664

Source: HPA Centre for Infections

** KC60 codes were created/revised in quarter 3, 2002

- data not available/applicable



England:

Number of new STI episodes seen at genitourinary medicine clinics by sex, 1998 - 2007

Diagnoses and/or treatment of infection or disease			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
A1,A2	Primary and secondary infectious syphilis	Male (all)	87	156	247	618	1,059	1,355	1,768	2,143	2,140	2,163
		-of which MSM	23	52	119	351	613	759	903	1,268	1,230	1,293
		Female	44	55	75	99	137	183	266	348	282	274
		Total	131	211	322	717	1,196	1,538	2,034	2,491	2,422	2,437
A3	Early latent syphilis (first 2 years)	Male (all)	87	122	126	268	330	425	526	660	669	807
		-of which MSM	18	30	48	114	177	194	271	384	410	462
		Female	58	75	93	99	121	150	197	175	172	151
		Total	145	197	219	367	451	575	723	835	841	958
A4,A5,A6	Other acquired syphilis	Male (all)	625	671	778	847	866	1,017	1,111	1,076	1,094	1,094
		-of which MSM	75	93	87	122	130	157	157	178	241	226
		Female	370	405	470	638	746	805	864	835	724	811
		Total	995	1,076	1,248	1,485	1,612	1,822	1,975	1,911	1,818	1,905
A7	Congenital syphilis, aged under 2	Male (all)	0	*	5	*	*	*	*	6	*	*
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	*	*	*	0	5	0	*	8	*	*
		Total	*	*	9	*	7	*	7	14	6	*
A8	Congenital syphilis, aged 2 or over	Male (all)	*	*	9	*	*	*	5	9	*	*
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	*	*	*	6	*	*	*	13	*	*
		Total	*	*	13	*	14	5	9	22	8	*
A9	Epidemiological treatment of suspected syphilis	Male (all)	19	24	26	61	157	255	366	486	654	734
		-of which MSM	0	7	6	23	81	148	210	283	413	474
		Female	36	34	26	30	49	66	89	122	125	159
		Total	55	58	52	91	206	321	455	608	779	893
B1,B2	Uncomplicated gonorrhoea	Male (all)	8,446	10,669	14,272	15,654	16,940	16,268	14,514	12,683	12,296	11,816
		-of which MSM	1,677	1,812	2,867	3,454	3,298	3,653	3,619	3,922	4,069	3,458
		Female	4,089	4,880	6,222	6,744	7,417	7,224	6,265	5,019	5,013	5,386
		Total	12,535	15,549	20,494	22,398	24,357	23,492	20,779	17,702	17,309	17,202
B3	Gonococcal ophthalmia neonatorum	Male (all)	*	5	7	7	8	7	10	9	20	12
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	5	7	6	9	9	*	14	81	8	14
		Total	8	12	13	16	17	*	24	90	28	26
B4	Epidemiological treatment of suspected gonorrhoea	Male (all)	1,421	1,557	2,112	2,588	3,012	3,301	3,449	3,516	3,723	4,217
		-of which MSM	441	429	597	815	879	1,080	1,103	1,235	1,336	1,372
		Female	1,519	1,789	2,189	2,563	2,944	3,134	3,064	3,075	2,928	3,263
		Total	2,940	3,346	4,301	5,151	5,956	6,435	6,513	6,591	6,651	7,480
B5	Complicated gonococcal infection - including PID and epididymitis	Male (all)	78	74	79	95	106	99	135	104	89	99
		-of which MSM	11	12	8	14	25	10	19	31	18	17
		Female	183	237	260	273	272	315	262	225	235	257
		Total	261	311	339	368	378	414	397	329	324	356

England: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
C1,C2,C3	Chancroid/LGV/Donovanosis	Male (all)	43	28	30	29	34	60	86	258	213	251
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	24	23	31	33	37	76	63	59	67	120
		Total	67	51	61	62	71	136	149	317	280	371
C4A,C4C	Uncomplicated genital chlamydial infection	Male (all)	18,937	21,776	26,579	29,562	34,331	37,996	42,681	45,155	48,974	53,468
		-of which MSM	454	621	962	1,365	1,422	1,725	1,789	2,260	2,802	3,109
		Female	24,975	29,184	34,791	38,618	43,786	47,520	50,267	50,775	50,306	53,691
		Total	43,912	50,960	61,370	68,180	78,117	85,516	92,948	95,930	99,280	107,159
C4B	Complicated genital chlamydial infection - including PID and epididymitis	Male (all)	331	432	528	545	571	637	653	723	778	802
		-of which MSM	31	25	23	35	32	52	45	70	68	63
		Female	1,521	1,763	2,114	2,074	2,134	2,091	2,311	2,236	2,170	2,526
		Total	1,852	2,195	2,642	2,619	2,705	2,728	2,964	2,959	2,948	3,328
C4D	Chlamydial ophthalmia neonatorum	Male (all)	15	15	19	15	15	18	42	15	13	12
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	15	12	23	28	25	25	34	22	19	5
		Total	30	27	42	43	40	43	76	37	32	17
C4E	Epidemiological treatment of suspected genital chlamydial infection	Male (all)	9,086	11,291	14,023	17,345	21,502	22,531	24,371	25,987	25,225	28,658
		-of which MSM	189	243	405	620	1,032	1,282	1,372	1,627	1,919	2,243
		Female	6,907	8,243	10,118	12,096	14,207	14,651	16,668	18,494	18,631	20,568
		Total	15,993	19,534	24,141	29,441	35,709	37,182	41,039	44,481	43,856	49,226
C4H**	Uncomplicated non-gonococcal/non-specific urethritis in males, or treatment of mucopurulent cervicitis in females	Male (all)	51,951	53,214	56,007	59,467	62,040	67,292	69,870	70,396	69,809	68,004
		-of which MSM	3,339	3,375	3,847	4,566	4,470	5,000	5,051	5,195	5,739	5,324
		Female	0	0	0	0	0	6,636	7,417	7,381	6,715	6,427
		Total	51,951	53,214	56,007	59,467	62,040	73,928	77,287	77,777	76,524	74,431
C4I	Epidemiological treatment of NSGI	Male (all)	5,168	6,053	6,141	5,907	5,899	5,958	6,213	5,747	5,615	6,045
		-of which MSM	442	720	608	674	605	734	791	738	818	768
		Female	18,133	18,504	18,773	19,558	18,534	16,915	17,097	16,924	15,964	16,249
		Total	23,301	24,557	24,914	25,465	24,433	22,873	23,310	22,671	21,579	22,294
C5	Complicated infection (non-chlamydial/non-gonococcal) - including PID and epididymitis	Male (all)	2,431	3,008	3,498	3,713	3,874	3,956	4,364	5,025	4,712	5,182
		-of which MSM	153	206	198	237	216	239	275	340	327	346
		Female	9,858	10,228	10,799	10,943	11,324	10,752	12,512	11,969	12,362	14,103
		Total	12,289	13,236	14,297	14,656	15,198	14,708	16,876	16,994	17,074	19,285
C6A	Trichomoniasis	Male (all)	298	641	731	733	568	285	297	330	311	386
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	5,877	5,448	6,036	6,279	6,628	6,224	5,924	5,058	5,177	5,401
		Total	6,175	6,089	6,767	7,012	7,196	6,509	6,221	5,388	5,488	5,787
C6B	Anaerobic/Bacterial vaginosis & anaerobic balanitis	Male (all)	1,403	1,388	1,314	1,392	1,391	1,444	1,420	1,474	1,640	1,637
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	51,381	55,518	60,667	64,884	70,909	71,917	69,426	70,642	73,559	85,311
		Total	52,784	56,906	61,981	66,276	72,300	73,361	70,846	72,116	75,199	86,948
C6C	Other vaginosis/vaginitis/balanitis	Male (all)	12,324	11,996	12,049	11,972	11,660	11,827	11,644	11,053	10,909	12,744
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	8,713	7,643	7,079	6,753	5,680	5,382	5,816	4,850	4,313	4,607
		Total	21,037	19,639	19,128	18,725	17,340	17,209	17,460	15,903	15,222	17,351

England: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	
C7A	Anogenital candidosis	Male (all)	7,523	7,219	6,958	7,165	6,828	5,970	5,947	5,591	6,375	7,572	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	63,016	64,196	65,686	69,918	70,401	66,661	62,919	59,340	58,357	65,240	
		Total	70,539	71,415	72,644	77,083	77,229	72,631	68,866	64,931	64,732	72,812	
C7B	Epidemiological treatment of C6 & C7	Male (all)	3,630	3,621	3,351	3,729	4,177	4,263	4,197	4,373	4,158	4,709	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	6,250	6,395	5,773	6,676	6,674	7,062	7,920	7,307	7,231	8,105	
		Total	9,880	10,016	9,124	10,405	10,851	11,325	12,117	11,680	11,389	12,814	
C8,C9	Scabies/pediculosis pubis	Male (all)	4,565	4,491	4,611	4,178	3,592	3,058	2,420	1,884	1,865	1,777	
		-of which MSM	762	741	826	805	708	597	501	448	468	426	
		Female	1,505	1,320	1,150	948	707	504	393	295	258	266	
		Total	6,070	5,811	5,761	5,126	4,299	3,562	2,813	2,179	2,123	2,043	
C10A	Anogenital herpes simplex – first attack	Male (all)	6,140	6,031	6,175	6,489	6,522	6,449	6,438	6,800	7,520	9,046	
		-of which MSM	301	338	384	399	479	468	458	550	526	624	
		Female	9,675	9,849	9,972	10,565	10,988	10,708	10,514	10,818	11,926	14,497	
		Total	15,815	15,880	16,147	17,054	17,510	17,157	16,952	17,618	19,446	23,543	
C10B	Anogenital herpes simplex – recurrence	Male (all)	5,560	5,983	6,095	6,204	6,376	6,373	6,072	6,250	6,630	7,172	
		-of which MSM	356	396	442	397	391	462	433	498	514	564	
		Female	6,798	7,226	7,345	8,049	8,142	8,009	7,724	8,013	8,374	9,683	
		Total	12,358	13,209	13,440	14,253	14,518	14,382	13,796	14,263	15,004	16,855	
C11A	Anogenital warts – first attack	Male (all)	30,782	31,846	31,975	32,700	33,987	34,642	36,418	36,368	37,479	40,328	
		-of which MSM	1,501	1,565	1,673	1,771	1,963	2,108	2,032	2,258	2,321	2,462	
		Female	28,899	29,311	28,686	29,723	29,951	30,637	31,799	32,333	33,253	36,340	
		Total	59,681	61,157	60,661	62,423	63,938	65,279	68,217	68,701	70,732	76,668	
C11B	Anogenital warts – recurrence	Male (all)	21,847	22,141	22,226	22,579	22,061	21,813	22,228	23,288	24,642	27,282	
		-of which MSM	1,228	1,211	1,213	1,376	1,387	1,548	1,615	1,544	1,729	2,004	
		Female	12,448	12,293	12,419	12,170	12,137	12,501	12,792	13,710	14,994	16,957	
		Total	34,295	34,434	34,645	34,749	34,198	34,314	35,020	36,998	39,636	44,239	
C11C	Anogenital warts – re-registered cases	Male (all)	8,942	9,314	9,047	8,988	8,464	8,803	8,499	8,778	9,102	9,294	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	7,408	7,597	7,219	7,148	6,517	6,616	6,450	6,810	6,773	6,778	
		Total	16,350	16,911	16,266	16,136	14,981	15,419	14,949	15,588	15,875	16,072	
C12	Molluscum contagiosum	Male (all)	4,363	4,763	5,100	5,571	6,187	6,940	8,045	8,846	9,623	10,541	
		-of which MSM	209	215	206	225	209	250	258	286	311	340	
		Female	2,373	2,637	2,691	3,003	3,250	3,485	3,892	4,239	4,507	4,863	
		Total	6,736	7,400	7,791	8,574	9,437	10,425	11,937	13,085	14,130	15,404	
C13	Antigen positive viral hepatitis B	Male (all)	551	527	635	704	931	0	0	0	0	0	
		-of which MSM	106	102	128	155	195	0	0	0	0	0	
		Female	203	249	230	352	472	0	0	0	0	0	
		Total	754	776	865	1,056	1,403	0	0	0	0	0	
C13A**	Viral hepatitis B (HbsAg positive): first diagnosis**	Male (all)	-	-	-	-	-	602	649	637	624	673	
		-of which MSM	-	-	-	-	-	111	88	90	60	60	
		Female	-	-	-	-	-	271	325	325	340	353	
		Total	-	-	-	-	-	873	974	962	964	1,026	

England: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
C13B**	**number of which were acute viral hepatitis B	Male (all)	-	-	-	-	-	46	54	44	40	40
		-of which MSM	-	-	-	-	-	14	19	14	7	7
		Female	-	-	-	-	-	10	12	25	26	17
		Total	-	-	-	-	-	56	66	69	66	57
C13C**	Viral hepatitis B: subsequent presentation	Male (all)	-	-	-	-	-	342	459	414	431	489
		-of which MSM	-	-	-	-	-	121	205	206	228	167
		Female	-	-	-	-	-	150	168	143	188	200
		Total	-	-	-	-	-	492	627	557	619	689
C14**	Viral hepatitis C: first diagnosis (previous to 2003 C14 was other viral hepatitis)	Male (all)	1,033	918	929	918	1,044	1,004	1,039	941	858	798
		-of which MSM	129	83	103	104	140	110	123	115	117	81
		Female	431	425	422	380	484	462	438	398	353	292
		Total	1,464	1,343	1,351	1,298	1,528	1,466	1,477	1,339	1,211	1,090
E1A	New HIV diagnosis - asymptomatic	Male (all)	1,104	1,063	1,184	1,655	2,302	2,656	2,769	2,852	2,678	3,310
		-of which MSM	546	518	590	740	737	1,094	1,201	1,295	1,223	1,377
		Female	351	573	670	1,109	1,977	2,106	2,054	2,036	1,730	2,200
		Total	1,455	1,636	1,854	2,764	4,279	4,762	4,823	4,888	4,408	5,510
E1B**	Asymptomatic HIV infection – subsequent presentation	Male (all)	6,165	6,505	7,588	8,375	10,058	0	0	0	0	0
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1,785	2,096	2,860	3,792	5,532	0	0	0	0	0
		Total	7,950	8,601	10,448	12,167	15,590	0	0	0	0	0
E1BE2B**	Subsequent HIV presentation (not AIDS)	Male (all)	-	-	-	-	-	25,773	30,935	37,629	39,162	39,653
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	-	-	-	-	-	12,209	16,402	20,525	21,543	23,011
		Total	-	-	-	-	-	37,982	47,337	58,154	60,705	62,664
E2A	New HIV diagnosis: symptomatic (not AIDS)	Male (all)	567	514	470	641	744	664	669	735	616	473
		-of which MSM	285	278	222	300	244	263	246	253	224	148
		Female	191	195	198	326	445	441	385	442	357	311
		Total	758	709	668	967	1,189	1,105	1,054	1,177	973	784
E2B**	HIV infection with symptoms, not AIDS – subsequent presentation	Male (all)	7,744	7,788	8,880	9,139	9,265	0	0	0	0	0
		-of which MSM	0	0	0	0	0	0	0	0	0	0
		Female	1,694	1,854	2,230	2,623	3,310	0	0	0	0	0
		Total	9,438	9,642	11,110	11,762	12,575	0	0	0	0	0
E3A**	AIDS – first presentation	Male (all)	427	377	360	430	436	0	0	0	0	0
		-of which MSM	188	151	178	167	155	0	0	0	0	0
		Female	127	110	139	214	227	0	0	0	0	0
		Total	554	487	499	644	663	0	0	0	0	0
E3A1**	AIDS: first presentation - new HIV diagnosis	Male (all)	-	-	-	-	-	176	217	226	189	225
		-of which MSM	-	-	-	-	-	69	61	76	64	74
		Female	-	-	-	-	-	137	116	150	89	95
		Total	-	-	-	-	-	313	333	376	278	320
E3A2**	AIDS: first presentation - HIV diagnosed previously	Male (all)	-	-	-	-	-	236	153	161	157	112
		-of which MSM	-	-	-	-	-	94	74	60	82	54
		Female	-	-	-	-	-	133	63	70	69	46
		Total	-	-	-	-	-	369	216	231	226	158

England: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	
E3B	AIDS – subsequent presentation	Male (all)	5,261	5,396	6,119	6,683	7,605	9,539	10,273	11,637	11,559	11,213	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	1,071	1,229	1,551	2,001	2,584	3,999	4,319	4,965	5,147	5,353	
		Total	6,332	6,625	7,670	8,684	10,189	13,538	14,592	16,602	16,706	16,566	
D2A	Urinary tract infection	Male (all)	1,927	2,040	1,910	1,981	2,522	2,483	2,479	2,028	1,987	2,258	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	8,711	9,064	9,351	9,431	9,152	8,717	9,772	9,351	9,678	12,041	
		Total	10,638	11,104	11,261	11,412	11,674	11,200	12,251	11,379	11,665	14,299	
D2B	Other conditions requiring treatment at GUM clinic	Male (all)	50,924	52,418	54,692	54,839	57,505	64,250	64,697	66,904	68,270	78,987	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	48,031	48,791	53,560	54,402	54,721	53,126	54,501	55,624	56,746	64,249	
		Total	98,955	101,209	108,252	109,241	112,226	117,376	119,198	122,528	125,016	143,236	
D3	Other episodes not requiring treatment	Male (all)	97,722	100,901	106,796	125,321	141,931	133,675	160,686	188,501	208,137	236,192	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	-
		Female	107,328	110,295	113,953	131,498	146,874	135,640	158,352	190,194	207,600	242,457	
		Total	205,050	211,196	220,749	256,819	288,805	269,315	319,038	378,695	415,737	478,649	
P1A	HIV antibody test (no sexual health screen)	Male (all)	71,699	73,635	83,890	109,903	145,623	48,164	42,841	39,940	33,954	31,934	
		-of which MSM	10,917	10,869	12,125	15,980	18,223	8,154	7,455	7,746	7,277	6,896	
		Female	57,599	59,051	67,569	92,977	129,933	37,024	30,537	29,874	25,348	24,551	
		Total	129,298	132,686	151,459	202,880	275,556	85,188	73,378	69,814	59,302	56,485	
P1B	HIV antibody test offered and refused	Male (all)	12,582	12,962	14,326	19,265	42,360	95,065	116,163	112,939	107,593	112,890	
		-of which MSM	1,089	1,396	1,137	1,643	3,102	5,937	6,237	6,278	6,286	6,240	
		Female	11,775	12,432	13,612	19,130	47,485	109,531	133,388	131,478	127,798	141,713	
		Total	24,357	25,394	27,938	38,395	89,845	204,596	249,551	244,417	235,391	254,603	
P2	Hepatitis B vaccination (1st dose only)	Male (all)	9,847	9,989	10,885	11,859	13,313	14,464	15,858	15,516	16,024	16,942	
		-of which MSM	4,835	4,707	5,218	5,850	6,604	7,838	8,348	8,432	8,375	8,560	
		Female	2,597	2,491	2,630	3,142	3,909	4,092	4,982	5,253	5,564	6,107	
		Total	12,444	12,480	13,515	15,001	17,222	18,556	20,840	20,769	21,588	23,049	
P3	Contraception (excluding condom provision)	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	31,801	30,263	32,771	33,646	31,444	32,774	33,858	34,563	37,969	45,490	
		Total	31,801	30,263	32,771	33,646	31,444	32,774	33,858	34,563	37,969	45,490	
P4A	Cervical cytology - minor abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-	
		-of which MSM	-	-	-	-	-	-	-	-	-	-	
		Female	9,387	7,583	6,214	4,744	3,663	5,212	3,351	1,676	1,362	1,389	
		Total	9,387	7,583	6,214	4,744	3,663	5,212	3,351	1,676	1,362	1,389	

England: Number of new STI episodes by sex, 1998 - 2007 continued

			1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
P4B	Cervical cytology - major abnormality	Male (all)	-	-	-	-	-	-	-	-	-	-
		-of which MSM	-	-	-	-	-	-	-	-	-	-
		Female	1,346	1,184	1,120	1,024	751	912	793	383	291	263
		Total	1,346	1,184	1,120	1,024	751	912	793	383	291	263
S1**	Sexual health screen (no HIV antibody test)	Male (all)	-	-	-	-	-	141,976	142,890	135,850	131,887	136,859
		-of which MSM	-	-	-	-	-	9,684	9,833	10,731	11,977	12,127
		Female	-	-	-	-	-	176,335	174,285	167,932	161,301	173,379
		Total	-	-	-	-	-	318,311	317,175	303,782	293,188	310,238
S2**	HIV antibody test and sexual health screen	Male (all)	-	-	-	-	-	158,950	208,016	255,017	298,276	354,585
		-of which MSM	-	-	-	-	-	15,882	18,964	22,812	27,644	31,092
		Female	-	-	-	-	-	156,028	201,819	251,341	292,024	357,978
		Total	-	-	-	-	-	314,978	409,835	506,358	590,300	712,563
Total new STI diagnoses		Male (all)	130,123	138,706	151,486	161,650	172,857	182,533	191,344	194,528	199,292	207,871
		-of which MSM	9,292	9,758	11,925	14,262	14,416	16,287	16,458	18,252	19,390	19,061
		Female	89,565	95,703	103,695	110,737	119,053	129,039	134,440	133,383	134,447	146,757
		Total	219,688	234,409	255,181	272,387	291,910	311,572	325,784	327,911	333,739	354,628
Total other STI diagnoses		Male (all)	72,376	77,248	83,561	89,441	96,061	106,053	114,278	125,254	128,272	136,308
		-of which MSM	2,937	3,280	3,584	4,308	4,837	5,699	6,026	6,547	7,462	8,167
		Female	58,386	61,487	65,469	71,024	75,098	79,232	85,802	93,842	95,478	103,056
		Total	130,762	138,735	149,030	160,465	171,159	185,285	200,080	219,096	223,750	239,364
Total other GUM clinic diagnoses		Male (all)	79,315	80,127	81,838	82,700	86,058	92,231	92,585	93,459	95,292	109,907
		-of which MSM	235	185	231	259	335	356	435	425	412	315
		Female	197,469	201,048	210,102	218,564	222,907	219,882	215,441	210,064	212,444	242,067
		Total	276,784	281,175	291,940	301,264	308,965	312,113	308,026	303,523	307,736	351,974
Total services provided		Male (all)	191,850	197,487	215,897	266,348	343,227	592,294	686,454	747,763	795,871	889,402
		-of which MSM	16,841	16,972	18,480	23,473	27,929	47,495	50,837	55,999	61,559	64,915
		Female	211,100	214,532	230,535	280,393	359,645	651,424	737,221	810,635	857,604	991,675
		Total	402,950	412,019	446,432	546,741	702,872	1,243,718	1,423,675	1,558,398	1,653,475	1,881,077

Source: HPA Centre for Infections

** KC60 codes were created/revised in quarter 3, 2002

- data not available/applicable

* confidential data

**Supplementary Tables of Sexually
Transmitted Infections in the United Kingdom**



Code Groups
New STI Diagnoses
Chlamydial infection (uncomplicated and complicated)
Gonorrhoea (uncomplicated and complicated)
Infectious syphilis
Genital Herpes simplex (first attack)
Genital warts (first attack)
New HIV diagnosis
Non-specific genital infection (uncomplicated and complicated)
Chancroid/lymphogranuloma venerum (LGV)/Donovanosis
Molluscum contagiosum
Trichomoniasis
Scabies
Pediculus pubis
Other STI Diagnoses
Early latent, congenital and other acquired syphilis
Recurrent genital Herpes simplex
Recurrent and re-registered genital warts
Subsequent HIV presentations (including AIDS)
Ophthalmia neonatorum (chlamydial or gonococcal)
Epidemiological treatment of suspected STIs (syphilis, chlamydia, gonorrhoea, non-specific genital infection)
Other diagnoses made at GUM clinics
Viral hepatitis B
Viral hepatitis C
Vaginosis and balanitis (including epidemiological treatment)
Anogenital candidiasis (including epidemiological treatment)
Urinary tract infection
Cervical abnormalities
Other conditions requiring treatment at a GUM clinic
Services provided
HIV antibody test
Sexual health screen
Hepatitis B vaccination
Contraception (excluding condom provision)
Other episode not requiring treatment